



GETTING THE BEST FROM ACELEPRYN

YOUR 7 POINT PLAN FOR APPLICATION ON GOLF COURSES AND RACE COURSES



1 Apply preventatively. Acelepryn is only active on 1st and 2nd instar. Larger grubs will not be controlled.



2 Apply to the turf during the period when peak numbers of target species are observed. Sites with a history of attack have a high probability to be re-infested.



3 Mow prior to application.



4 Apply at 0.6L/ha, in a minimum of 600 litres of water, using a Syngenta XC soil nozzle or suitable air induction 08 nozzle between 2–4 bar pressure.



5 Where ever possible irrigate in with 3-6 mm water after application, avoiding run-off. If irrigation is not possible then natural rainfall will be required to move the product down through the organic matter layer.



6 Apply in tank mix or sequence with a wetting agent where significant organic matter is present or if there are hydrophobic areas. Syngenta wetting agent Qualibra is compatible with Acelepryn.



7 Delay mowing after application until irrigation has taken place or rainfall has occurred. If Acelepryn has not been washed in then delay mowing for as long as possible and return the clippings if feasible.

RESEARCH HAS SHOWN ACELEPRYN CAN PROVIDE ON AVERAGE 90% CONTROL WITH A WELL PLANNED AND EXECUTED APPLICATION

HIGH AND MIXED POPULATIONS WILL NEED A LONG TERM PROGRAMMED APPROACH