

GETTING THE BEST FROM ACELEPRYN®

YOUR 7 POINT PLAN FOR APPLICATION



1 Apply preventatively. Acelepryn is only effective on 1st and 2nd instar. Larger grubs / larvae will not be controlled.



4 Apply at 0.6L/ha, in 500-600 litres of water, using a Syngenta XC soil nozzle or suitable air induction 08 nozzle between 2-4 bar pressure.



2 Chafers: Apply once peak flight has been observed (typically June).
Leatherjackets: Adult crane fly activity may not peak until mid-October. Always check peak flight has occurred for application timing.



5 Ensure the soil profile is not hydrophobic at the time of application. Syngenta wetting agent Qualibra is compatible with Acelepryn for convenient application in tank-mix or sequence.

Sites with historic high pressure have increased probability of re-infestation.
Follow reports of adult activity and record sightings on PestTracker at:
www.syngentaturf.co.uk/turf-pest-tracker



6 Keep surfaces irrigated where possible, to attract pest larvae to towards the surface.



3 Mow prior to application.



7 Delay mowing for as long as possible and return the clippings if feasible.

USE ACELEPRYN AS PART OF AN ITM PROGRAMME, INCLUDING CULTURAL PRACTICES AND COMPLEMENTARY NEMATRIDENT BENEFICIAL NEMATODE TREATMENTS.

HIGH AND MIXED POPULATIONS WILL NEED A LONG TERM PROGRAMMED APPROACH

Follow [#PestTracker](https://twitter.com/SyngentaTurfUK) for timing information support and [@SyngentaTurfUK](https://twitter.com/SyngentaTurfUK) for turf management advice.
For technical support contact product.technical_enquiries@syngenta.com or 0800 1696058.

 **Acelepryn®**

syngenta.